

A11100 988208

NAT'L INST OF STANDARDS & TECH R.I.C.



A11100988208


Spencer, L. V. (Low/Structure shielding
QC100 .U57 V570:1980 C.1 NBS-PUB-C 1980



NBS SPECIAL PUBLICATION 570

U.S. DEPARTMENT OF COMMERCE/National Bureau of Standards


Structure Shielding Against Fallout Gamma Rays From Nuclear Detonations




AUTHENTICATED
U.S. GOVERNMENT
INFORMATION

GPO







I.4 The multi-slab concrete sandwich for the experiments on penetration of fallout gamma rays, carried out by F. Titus as part of the Plumbob test series [39].



II.1 Yield data for several cases of nuclear fission [2].



II.4 Principle contributions to the activity of unfractionated fission products from thermonuclear neutron fission of ^{238}U [28].



II.17 Curves give attenuation factor versus concrete thickness for a smooth isotropic plane source emitting various fission spectra [93]. The heavy plus signs lying above the curves give experimental data due to Titus (shot B, 2 hours after detonation) [97].

throughout the cloud. The cloud is subdivided into segments, and each segment is repeatedly assigned a specified particle size and called a wafer. These are transported according to wind patterns and estimated fall rates.

The prototype of the scaling studies was due to C. F. Ksanda and coworkers in 1953 [113] at the Naval Radiological Defense Laboratory (NRDL), whose study was based on fallout patterns from Operation Jangle (1951) which were described by R. K. Laurino and I. B. Poppoff [114]. Results of early scaling studies were incorporated in the 1957 edition of Effects of Nuclear Weapons [12]; but W. W. Kellogg, in his 1957 testimony to Congress, documented the discouragingly poor agreement among different fallout prediction systems at that time [115]. A fallout symposium held in March 1957 concluded that "without a more thorough understanding of the mechanics of the fallout process, from the early stages of the fireball until a major fraction is deposited on the ground, we cannot expect to construct a fallout model that will yield reasonably accurate predictions" [116].

Much early work on fallout models was incompletely published, partially classified, or suited only to some highly specialized application. The 1957 Congressional Hearings are a good source for early results and general approaches, particularly Kellogg's presentation [115], and Schuert's discussion [117]. Unclassified and reasonably complete presentations of particular methods a few years later include A. D. Anderson's paper in 1961 [116] and the 1960 report by Callahan and coworkers [118].

Widespread work on fallout prediction systems in the late 1950's led to a second symposium in September, 1962, at which intercomparisons of many models were discussed on the basis of hypothetical burst conditions previously circulated to interested groups. This conference was fortunately followed up by extensive analysis of 18 of the models, even though many models were still not well documented [119,120]. Comparisons again showed that predictions of different models were very diverse.

A third symposium was held in April, 1966, and was far better focussed on the basic physics problems fundamental to the important processes [121]. By this time it was clear that only a few models would receive extensive additional development and use. In fact, much of the symposium consisted of reports about work on subunits of the DELFIC prediction system, a wafer system intended to satisfy requirements by all the different branches of the Department of Defense, thus superceding a large fraction of the existing models.

In the remaining paragraphs of this section we sketch four systems for calculating fallout intensities and spectra which remain relevant to current applications. These are the WSEG model, a Miller model, the DELFIC model, and a SEER model.

One should be aware throughout this discussion that the intercomparisons of reference [120] have not been fully extended to the more recent models or versions of older models.²¹ Further, no systematic study seems to have been made of any model in comparison with data from most of the tests which have been conducted. This is largely because of the state of information about the test detonations, which is fragmentary except for a few cases. But there is also a difficulty of interpretation of differences because of the large number of variables involved.

This problem of pinning prediction systems to adequate, incontrovertible data leads to the opinion that these systems have their main application in hypothetical attack studies, and that they would not be dependable as a sole basis for fallout intensity estimates after a nuclear attack or for operational decisions on which the lives of people near a detonation might depend.

²¹A step in this direction was taken by C. J. Seery [122], in a paper describing a "TINCAN" model suggested by I. Russell. Sensitivity studies were included, with data on cloud radius, cloud height, particle fall rate, refractory mass fraction, and particle frequency distribution. Comparisons of TINCAN with the other models described here were made.

a. The WSEG Model²²

Of the scaling models, that of G. E. Pugh and R. J. Galiano is the currently most important because it is used nearly universally in the hypothetical attack studies which attempt to assess damage to this country [119]. In this model the cloud activity is described by a Gaussian distribution whose height and lateral spread are functions of the yield. Neither particle size distribution nor fall rates are explicitly used in this model; and the cloud is not divided into wafers. Instead, there is a carefully specified function of the time after detonation which determines the rate of activity deposition on the ground. A scaling rule changes "time" in this function to downwind distance, by use of an effective wind speed. A crosswind shear dependent on cloud thickness must also be specified.

Dose rate on the ground is specified by the product of separate functions of the downwind and crosswind distances: The crosswind function is a Gaussian, and the downwind function is the product of a falling exponential function of downwind distance with a cumulative normal function.

This model is implemented by a very fast code, clearly suitable to studies which must cumulate the effect of many bursts, when details tend to be unimportant because of effects of superposition.

b. The Miller Model

A scaling approach was developed by Miller in the period 1962-64 based on data from the test shots Jangle 5 and Castle Bravo, and has been described in a 1969 paper [124]. More complex and physical than the WSEG model, it has been used in some damage assessment studies.²³

²²For Weapons System Evaluation Group [123].

²³A later model (see ref. [125] by C. F. Miller is more empirical and comparable with DELFIC. While Miller's most recent model reflects the state of the art, we prefer to describe here Miller's earlier, more conceptual approach.

The Miller model distinguishes between the stem and the cloud of the mushroom due to a ground burst. The stem generates a high intensity "ridge" downwind, with intensity decreasing laterally. Superimposed on this is the broader cloud fallout pattern carried by the wind to greater distances because of its initially greater altitude. For both stem and cloud, intensities and distances are specified along the pattern centerline as functions of the yield and the wind velocity, and along a line of maximum lateral extent by use also of wind correction and shear factors. Both stem and cloud patterns are then described by contours in the shape of half-ellipses smoothly joined together and in agreement with the centerline and width values. The two components are summed to get the total "intensity", which is identified as a specific detector measurement 3 ft above an extended open area. The result in a simple wind pattern is to reproduce the fallout as a sort of shadow of the mushroom cloud, a characteristic observed with atmospheric nuclear tests.

Particle fall from the stem is assumed to be unaffected by lateral wind shear, but shear is taken into account in a simple way in the description of fallout from the cloud. Wind velocity is assumed to have an average magnitude at all altitudes. Particle concentrations are assumed to be uniform in the original mushroom cloud; but larger particles are assumed to fall from lower altitudes in the stem, and particle concentrations in the stem are assumed to decrease gradually with height. Contributions to exposure are functions of particle size expressed through fall speed; and at any location, the fallout varies between minimum and maximum fall speeds characteristic of the position. This leads to definite times of arrival and cessation of the fallout, and to exoisyre rate as a function of time and position, taking account of fractionation.

c. The DELFIC Model²⁴

Beginning about 1965, an effort was instituted by the Defense Atomic Support Agency (now the Defense Nuclear Agency) to develop a single model for all applications. It was intended that this model, called DELFIC, would combine the best features available. But the model would have to be a compromise in many respects and in this regard one more among many systems.

DELFIG consists of five principal modules: Initial Conditions, Cloud Rise, Transport, Particle-activity, and Output-processor.

The "Initial Conditions" module requires yield and height data, together with data on the type²⁵ and natural particle-size distribution of the soil. It identifies an effective starting time for cloud rise and specifies for that time the average temperature of the nuclear cloud, the total amount of entrained soil material together with the fractions in vapor and condensed phases, and the size-frequency distribution of the soil particles. In constructing the code, it was eventually concluded that the natural size distribution of soil particles could be used except possibly for large-yield detonations. Most other output data are determined by scaling relationships based on test measurements and on estimates of energy partition among different processes.

The "Cloud Rise" module is based on a modified version of a model developed by I. O. Huebsch [128-130]. There are two stages of the calculation. In the first, the cloud is treated as a whole, with uniform density and temperature, water and water vapor conditions, and turbulent kinetic energy density. Mass of the cloud can change due to entrainment and particle fallout and these affect the dynamic behavior. Vertical wind shear is accounted to increase entrainment

²⁴Department of DEdefense Land and Fallout Interpretive Code. The Monterey Symposium Report [126] has been superseded by [127] and should be used with caution.

²⁵In general, soil types can be classified as either siliceous (melting temperature near 1673 K) or calcareous (melting temperature near 2887 K).

through changes in shape which are not otherwise considered. Cloud volume and mass are related through assumption of an expanding oblate spheroidal shape.

In the second part of the Cloud Rise module, particle wafers are identified at the initial time by subdividing the cloud into horizontal disk segments and assigning to each segment a set of wafers, each with characteristic particle size interval. The wafers are then subdivided into horizontal segments to permit horizontal distortion; the upper and lower wafer boundaries are followed separately to take account of vertical stretching. The full set is then followed in space through the rise processes, as position and shape are modified by the forces effective. The resulting output is a set of distorted wafers distributed vertically and suitable for the transport process. Wafers with different particle size intervals will be at different heights after cloud rise.

The "Atmospheric Transport" module utilizes an atmosphere divided into sub-volumes, each with properties uniform throughout. Large cells are used to take account of region of macro-meteorological phenomena, while small cells can be used to take account of local circulation systems such as those affected by topographical features such as a mountain. With this approach, wind-patterns characteristic of certain regions and having considerably complexity, including vertical components, can be utilized. Standard settling rates are used based on formulae for spheres of different sizes. Wind effects and fallout during cloud rise are taken into account.

The "Particle Activity" module distributes activity with particle size according to a modified version of Freiling's radial-distribution model; that is to say, volatile chains are assumed to condense on surfaces while refractory chains are distributed through particle volumes, as previously discussed. Fractionation is thus taken into account and predictions for specific types of

nuclides can be made as well as for total activity. Induced activities are also included. Dose rates 3 ft above a smooth plane are determined according to the local spectrum.

The "Output Processor" module sets up the format for the various selections of output data of greatest interest [131].

d. SEER Models²⁶

While DELFIC was developed to be as realistic as the state-of-the-art permitted, this same requirement implies long and comparatively expensive runs. Other requirements for fast and cheap computations led to attempts to develop simplified versions of DELFIC, in competition with the WSEG Model but hopefully giving results more nearly equivalent to those of DELFIC for the intended applications. The SEER Models (Simplified Estimation of Exposure to Radiation) are of this type, with running times for a single case 1/50 to 1/100 that of DELFIC, and with rather good agreement for sample cases.

To achieve short running times assumptions such as the following are used: Starting point at stabilized cloud, with parameters from DELFIC-generated library, wind profiles constant in time and space, a log-normal particle size distribution, precalculation of falling times for deposition of cloud wafers, uniform cloud, and use of standard function types to represent lateral distributions relative to the "hot-line" for a cloud layer.

2. Continuing Problems of Fallout Phenomena

Research into different aspects of fallout phenomena has been at a low level for many years. This is due in part to the lack of new data; but much research has been cut back despite the possibility of significant progress. Hence we sketch here briefly the status of some representative problem areas. As our model we use a similar discussion in ref. [133], which was divided into the four categories which we use below. See also [13], Appendix VII.

²⁶See reference [132].

- [77] LaRiviere, P. D., Yu, S., and Miller, C. F., Intensity--Activity Relations for Shot Small-Boy, Proc. Fallout Phenomena Symposium, April 12-14, 1966, Monterey, Calif., 403-437 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [78] Freiling, E. C., Fractionation III. Estimation of Degree of Fractionation and Radionuclide Partition for Nuclear Debris, Report No. USNRDL-TR-680, 35 pages (September 12, 1963).
- [79] Nathans, M. W., Thews, R., Holland, W. D., and Benson, P. A., Particle Size Distribution in Clouds from Nuclear Airbursts, J. Geophysical Research 75, 7559-7572 (1970). Note also Freiling, E. C., A Comparison of the Fallout Mass-Size Distributions Calculated by Lognormal and Power-Law Models, Report No. USNRDL-TR-1105, (1967).
- [80] Cassidy, S. H., FORTRAN Statement of the Generalized Land Fallout Model Computer Program (GLFMCP), Report No. USNRDL-LR-176 (1966); and The Computer Program (TSABND) which Produces the Tables of Mass Chain Refractory and Fission Product Radioactivity Content for the Particle Activity Module of the DOD Fallout Model, Report No. USNRDL-LR-173 (1966).
- [81] Freiling, E. C., Kay, M. A., and Sanderson, J. V., Fractionation IV. Illustrative Calculations of the Effect of Radionuclide Fractionation on Exposure Dose Rate from Local Fallout, Report No. USNRDL-TR-715, 38 pages (Jan 6, 1964).
- [82] Cassidy, S. H., and Crocker, G. R., Sensitivity Analysis of the Radial Distribution Model. Sensitivity to Carrier Material, Yield, and Type of Fission, Report No. USNRDL-TR-67-70, 68 pages (May 22, 1967).
- [83] Tompkins, R. C., Department of Defense Land Fallout Prediction System, Vol. V--Particle Activity, Report No. NDL-TR-102, 264 pages (US Army Nuclear Defense Lab., Edgewood Arsenal, Md., Feb. 1968).
- [84] Cook, C. S., Energy Spectrum of Gamma Radiation from Fallout, Health Physics 4, 42-51 (1960).
- [85] Miller, C. F., Gamma Decay of Fission Products from the Slow Neutron Fission of ^{235}U , Report No. USNRDL-TR-187 (July 11, 1957).
- [86] Mather, R. L., Johnson, R. F., and Tomnovec, F. M., γ -Radiation Field Above Fallout-Contaminated Ground, Health Physics 8, 245-260 (1962).
- [87] LaRiviere, P. D., Estimating the Importance of Individual Radionuclides in Fission and Induced Product Mixtures, Report, SRI Project No. MU-5779, 10 pages (Stanford Research Institute, Menlo Park, Calif., June 1966).
- [88] Mather, R. L., Residual Gamma Radiation Fields From Induced Soil Activities, Report No. USNRDL-TR-240, 27 pages (April 9, 1958).
- [89] Klement, A. W., Jr., A Review of Potential Radionuclides Produced in Weapons Detonations, AEC Report WASH-1024, 102 pages (July 30, 1959).

- [90] Maerker, R. E., and Muckenthaler, F. J., Gamma-Ray Spectra Arising from Thermal-Neutron Capture in Elements Found in Soils, Concretes, and Structural Materials, Report No. ORNL-4382, 264 pages (Oak Ridge Natl. Lab., Oak Ridge, Tenn. 37830, August, 1969).
- [91] Groshev, L. V., Demidov, A. M., Lutsenko, V. N., and Pelekhov, V. I., Atlas of γ -Ray Spectra from Radiative Capture of Thermal Neutrons, 198 pages, translated from the Russian by J. B. Sykes, (Pergamon Press, 1959).
- [92] Knapp, H. A., Magnitude and Distribution of Weapon Effects for the Design of Shelters for Protection against Fallout, Research Paper P-194, 124 pages (Inst. for Defense Analyses, Weapons System Evaluation Division, Arlington, Va., July 1965); and USAEC External Gamma Doses and Dose Rates from the Fallout from Nuclear Explosions, Hearings on Civil Defense before a Subcommittee of the Committee on Government Operations, 86th Congress, p. 527 (May 16, 1960).
- [93] Tomnovec, F. M., and Ferguson, J. M., Neutron-Field and Induced-Activity Measurements -- Operation BREN, Report No. CEX-62.50, 42 pages (U.S. Atomic Energy Commission, Sept. 1965).
- [94] Dunning, G. M., Cobalt-60 Bombs, Health Physics 4, 52-54 (1960).
- [95] Russell, R. S., Bartlett, B. O., and Bruce, R. S., The Significance of Long-Lived Nuclides After a Nuclear War, AEC Symp. Series 24, 548-565 (U.S. Atomic Energy Commission, Dec. 1971).
- [96] Long-Term Worldwide Effects of Multiple Nuclear-Weapons Detonations, NAS-NRC Report, 213 pages (Natl. Acad. of Sciences, Washington, D.C., 1975).
- [97] Titus, F., Penetration in Concrete of Gamma Radiation from Fallout, NBS Report No. 6143, 38 pages (Natl. Bureau of Standards, Washington, D.C., Sept. 4, 1958).
- [98] Donovan, L. K., and Chilton, A. B., Dose Attenuation Factors for Concrete Slab Shields Covered with Fallout as a Function of Time after Fission, Tech. Report No. R-137, 31 pages (U.S. Naval Civil Eng. Lab., Port Hueneme, Calif., June 1, 1961).
- [99] Spencer, L. V., Structure Shielding Against Fallout Radiation from Nuclear Weapons, NBS Monograph 42, 92 pages (Natl. Bureau of Standards, Washington, D.C., June 1, 1962).
- [100] Cialella, C. M., et al., Shielding Effectiveness of Enclosure Shields in a Fallout Field, POR-2221 (1963). (Confidential)
- [101] French, R. L., Price, J. H., and Tompkins, K. W., An Analysis of the Performance of Enclosure Shields in a Real Fallout Field, Report No. RRA-T55, 110 pages (Radiation Research Associates, Inc., Ft. Worth, Texas, Dec. 31, 1965).
- [102] Cialella, C. M., and King, N. D., The Shielding Properties of Simple Enclosure Shields, Report No. BRL MR 1663 (Ballistic Research Laboratories, Aberdeen Proving Ground, Md., 1965).

- [103] Tomnovec, F. M., Ferguson, J. M., and Weldon, D. M., The Effect of a Changing Gamma-Ray Fallout Spectrum on the Ground Roughness Factor, Report No. USNRDL-TR-915, 30 pages (Sept. 22, 1965).
- [104] Huddleston, C. M., Klingler, Q. G., Burson, Z. G., and Kinkaid, R. M., Ground Roughness Effects on the Energy and Angular Distribution of Gamma Radiation from Fallout, Health Physics 11, 537-548 (1965).
- [105] Hogan, O. H., Zigman, P. E., and Mackin, J. L., Beta Spectra II. Spectra of Individual Negatron Emitters, Report No. USNRDL-TR-802, 212 pages (Dec. 16, 1964).
- [106] Boyd, A. E., and Morris, E. E., Spatial Distribution of Energy Dissipated by Fallout β -Rays, Health Physics 2, 321-325 (1960).
- [107] Heller, R. B., Energy and Time Beta Ray Spectra of Fission Products of U^{235} by Fission Neutrons and U^{238} by 14 MeV Neutrons, WSEG Research Memo. No. 19, 56 pages (Weapons System Evaluation Group, Feb. 16, 1961).
- [108] Hogan, O. L., Beta Spectra. III. Spectra of U^{235} Fission Products at Selected Times After Fission, Report No. USNRDL-TR-1093, 50 pages (Oct. 3, 1966).
- [109] Curtis, B. G., and Petty, J. S., A Program to Compute Beta Radiation Dosage, Vol. 1 - Program Description, Report No. ARC 67-48/TRC-67-20, 123 pages, (American Research Corp., Fullerton, Calif., Aug. 1, 1967).
- [110] Mackin, J. L., Brown, S. L., and Lane, W. B., Beta Radiation Dosimetry for Fallout Exposure Estimates: Comparison of Theory and Experiment, Report No. TRC-69-26, 64 pages (Stanford Research Institute, Menlo Park, Calif., June 1969).
- [111] Kutcher, J. W., and Wyman, M. E., An Experimental Study of the Time Dependence of the Beta Energy Spectrum from ^{235}U Fission Fragments, Nucl. Sci. Eng. 26, 435-446 (1966).
- [112] Mikhail, S. Z., Beta-Radiation Doses from Fallout Particles Deposited on the Skin, Report No. ESA-TR-71-01, 91 pages (Environmental Science Associates, Burlingame, Calif. 94010, Jan. 1971).
- [113] Ksanda, D. V., Minvielle, L., and Moskin, A., Scaling of Contamination Patterns, Surface and Underground Detonations, Report No. USNRDL-TR-1 (Classified Report) (1953).
- [114] Laurino, R. K., and Poppoff, I. G., Contamination Patterns at Operation JANGLE, Report No. USNRDL-399 (Classified Report) (1953).
- [115] Kellogg, W. W., The Nature of Radioactive Fallout and its Effect on Man, Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, May 27-29 and June 3, 1957, Part 1, 85'th Congress, 104-141 (U.S. Govt. Printing Office, Washington, D.C., 1957).
- [116] Anderson, A. D., A Theory for Close-In Fallout From Land-Surface Nuclear Bursts, J. Meteorology 18, 431 (August 1961).


- [117] Schuert, E. A., The Nature of Radioactive Fallout and its Effect on Man, Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, May 27-29 and June 3, 1957, Part 1, 85'th Congress, 267-307 (U.S. Govt. Printing Office, Washington, D.C., 1957).
- [118] Callahan, E., Rosenblum, L., Kaplan, J., and Batten, D., The Probable Fallout Threat over the Continental United States, Report No. TO-B 60-13, 198 pages (1960); see also Brooks, F. C., Callahan, E. D., Clark, E. T., Batter, J. F., and Kaplan, A. L., Radiological Defense Planning Guide, Report No. TOI 58-26, 327 pages (Technical Operations, Inc., Burlington, Mass. July 31, 1958).
- [119] Polan, M., An Analysis of the Fallout Prediction Models. Vol. 1: Analysis, Comparison, and Classification of Models, Report No. USNRDL-TRC-68, 145 pages (Sept. 8, 1966).
- [120] Seery, C. J., and Polan, M., An Analysis of the Fallout Prediction Models, Vol. II: Analysis, Comparison, and Evaluation of Model Predictions, Report No. NRDL-TRC-68-59, 74 pages (Aug. 28, 1968).
- [121] Fallout Phenomena Symposium, Proceedings, Part 1, April 12-14, 1966, Monterey, Calif., 486 pages (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [122] Seery, C. J., Fallout Model Analysis, Report No. TRC-68-46, 166 pages (Office of Civil Defense, Washington, D.C., Nov. 1968).
- [123] Pugh, G. E., and Galiano, R. J., An Analytical Model of Close-In Deposition of Fallout for Use in Operational-Type Studies, Research Memo. No. 10 (Weapons System Evaluation Group, 1959).
- [124] Miller, C. F., Biological and Radiological Effects of Fallout from Nuclear Explosions. Chapter 3. Distribution of Local Fallout, Report No. URS 702-1/TRC-68-61, 97 pages (URS Research Co., Burlingame, Calif. 94010, May 1969).
- [125] Miller, C. F. The Analysis and Correlation of Fallout Pattern Data. Part Two: Derivation of a Statistical Fallout Pattern Computational System, Dikewood Corp. Report No. DC-FR-1216-II, 100 pages, May 1973; Part Three: Computer Program User's Instructions, Dikewood Corp. Report No. DC-FR-1216-III, March 1974. Also, see Fallout Modifications Due to Unusual Burst Conditions, Dikewood Corp. Report No. DC-FR-1219, 66 pages, Dec. 1974; and Constraints of Prior Dose on Postattack Operational Planning for Areas Affected by Radioactive Fallout, Center for Planning and Research, Inc. Report, Aug. 31, 1975.
- [126] Tompkins, R. C., The DOD Land Fallout Prediction System, Fallout Phenomena Symposium, Proc., Part I, April 12-14, 1966, Monterey, Calif., 291-300 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).

- [127] Department of Defense Land Fallout Prediction System, Report No. DASA-1800, Vol. I, System Description, 24 pages (June 1966); Vol. II, Initial Conditions, 64 pages (Sept. 1966); Vol. III, Cloud Rise, 210 pages (Sept. 1970); Vol. IV, Atmospheric Transport, 306 pages (Feb. 1967); Vol. V, Particle Activity, 264 pages (Feb. 1968); Vol. VI, Output Processor, 112 pages (Feb. 1967); and Vol. VII, Operator's Manual 434 pages (Defense Atomic Support Agency, Washinton, D.C., April 1968).
- [128] Huebsch, I. O., Development of a Land Surface Burst Cloud Rise Model, Report No. USNRDL-LR-145 (November 1965).
- [129] Huebsch, I. O., The Energy Cycle and the Energy Cascade of the Atomic Cloud, Fallout Phenomena Symposium, Proc., Part I, April 12-14, 1966, Monterey, Calif., 129-180 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [130] Huebsch, I. O., Wind Shear, Turbulence, and Interface Criteria for Nuclear-Explosion Cloud, Debris and Fallout Models, Report No. NRDL-TR-69-72 (July 10, 1969).
- [131] Schwenke, T. W., Transport and Output Processor Modules for the DOD Fallout Prediction System, Fallout Phenomena Symposium, Proceedings, Part I, April 12-14, 1966, Monterey, Calif., 215-235 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [132] Lee, H., Wong, P. W., and Brown, S. L., Seer II: A New Damage Assessment Fallout Model, Report No. DNA 3008F, 137 pages (Defense Nuclear Agency, Washington D.C. 20305, May, 1972). AD 754144.
- [133] Fallout Phenomena Symposium, Proceedings, Part I, April 12-14, 1966, Monterey, Calif., Summaries and Recommendations of the Working Groups: Radiation Fields, 467-474 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [134] Heft, R. E., Phillips, W., and Steele, W., Radionuclide Distribution in the Particle Population Produced by the Schooner Cratering Detonation, Nucl. Tech. 11, 413-443 (July 1971).
- [135] Tompkins, R. C., Effect of Depth of Burial on Fallout from Atomic Demolition Munitions, Report No. BRL MR 2317 (Ballistic Research Lab., Aberdeen Proving Ground, Md., August 1973).
- [136] Knox, J. B., Prediction of Fallout from Subsurface Nuclear Detonations, AEC Symp. Series 5, 331-353 (U.S. Atomic Energy Commission, Nov. 1965).
- [137] A Precipitation Scavenging Model for Studies of Tactical Nuclear Operations, Vol. I, Theory and Preliminary Results, Report No. DNA 3661F-1, 61 pages (Mt. Auburn Research Associates, Inc. June 18, 1965). Note also observational data in the papers by Huff, F. A.; Saucier, W. J., et al; Hall, S. J., Gatz, D. F., and Dingle, A. N.; Kruger, P., et al; Kuroda, P. K., et al; and Volchok, H. L.; AEC Symp. Series 5, 507-628 (U.S. Atomic Energy Commission, November 1965).

- [138] Survival of Food Crops and Livestock in Event of Nuclear War, AEC Symp. Series 24, 718 pages (Brookhaven National Lab., Dec. 1971).
- [139] Radioecological Concentration Processes, Proc. International Symp., April 25-29, 1966, Stockholm, Sweden, 1040 pages (Pergamon Press, 1966).
- [140] Cowan, F. P., and Meinhold, C. B., Decontamination, Chapter 10, Survival and the Bomb, Methods of Civil Defense, 307 pages (Indiana U. Press, Bloomington, Indiana, 1969).
- [141] Straub, C. P., Decontamination of Food and Water Following Nuclear Attack, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 171-180 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April 1968).
- [142] Tamura, T., Movement of Cesium-137 by Runoff, Erosion, and Infiltration from a Soil Under Different Cover Conditions, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 149-168 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April 1968).
- [143] Mackin, J. L., and Mikhail, S. K., A Discussion of the State of the Art in Fallout Research, Reviews and Lectures No. 157, 73 pages (U.S. Naval Radiological Defense Lab., San Francisco, Calif., Dec. 7, 1965).
- [144] Nishiwaki, Y., Kawai, H., Hyono, A., Kondo, M., Goshi, N., Nishiwaki, J., Yamatera, H., Kudo, I., Mori, M., Azuma, T., Tsumori, K., and Nagayama, O., Studies on the Radioactivity of Bikini Ash, Research in the Effects and Influences of the Nuclear Bomb Test Explosions, page 461 (Japan Society for the Promotion of Science, Ueno, Tokyo, 1956).
- [145] Miller, C. F., Operation Ceniza-Arena: The Retention of Fallout Particles from Volcan Irazu (Costa Rica) by Plants and People, Part 2 247 pages; Part 2 Appendices, 281 pages (Dec. 1966); and Part 3, 255 pages, SRI Report No. TRC-68-17 (Dec. 1967); (Stanford Research Institute, Menlo Park, Calif.)
- [146] Evans, E. C., and Young, C. A., Differences in Transport and Deposition of Close-in and Intermediate Fallout, Fallout Phenomena Symposium, Proc., Part I, April 12-14, 1966, Monterey, Calif., 236-290 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [147] Soule, R. R., Gamma Ray Fields Above Rough Contaminated Surfaces, Fallout Phenomena Symposium, Proc., Part I, April 12-14, 1966, Monterey, Calif., 319-336 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [148] Knapp, H. A., Gamma Ray Exposure Dose from the Surface Deposition of Nuclear Test Fallout, 40 pages (U.S. Atomic Energy Commission, Div. of Biology and Medicine, Washington, D.C., July 21, 1962).
- [149] Radiological Factors Affecting Decision-Making in a Nuclear Attack, NCRP Report No. 42, 66 pages (National Council on Radiation Protection Measurements, Nov. 15, 1974).

- [150] Brown, S. L., Disintegration Rate Multipliers in Beta Emitter Dose Calculations, Fallout Phenomena Symposium, Proc., Part I, April 12-14, 1966, Monterey, Calif., 337-356 (Office of Civil Defense or Defense Atomic Support Agency, Washington, D.C., 1966).
- [151] Brown, S. L., and Yu, O. S., Computational Techniques in Beta Dose Problems, Report No. TRC-68-13 (Stanford Research Inst., Menlo Park, Calif., Jan. 1968).
- [152] Dunham, C. L., The Long-Range Biological Effects on Man, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 181-190 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April, 1968).
- [153] Brown, S., Lee, H., and Yu, O., Postattack Food Production and Water Contamination, SRI Report to OCD, 115 pages (Stanford Research Institute, Menlo Park, Calif., June 1968).
- [154] Radioactive Fallout from Nuclear Weapons Tests, AEC Symp. Series 5, 953 pages, (U.S. Atomic Energy Commission, Nov. 1965).
- [155] The Nature of Radioactive Fallout and its Effects on Man, Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, May 27-29 and June 3, 1957, Parts 1 & 2, 85'th Congress, 2065 pages (U.S. Govt. Printing Office, Washington, D.C., 1957).
- [156] Biological and Environmental Effects of Nuclear War, Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, June 22-26, 1959, Part 1, 87'th Congress, 966 pages (U.S. Govt. Printing Office, Washington, D.C., 1959).
- [157] Cole, R., Assessment of the Hazards of Fallout Radioiodines, Report for DCPA (Environmental Science Associates, Burlingame, Calif., Aug. 1972).
- [158] Damage to Livestock from Radioactive Fallout in the Event of Nuclear War, NAS-NRC Publication 1078, 93 pages (National Academy of Sciences, National Research Council, Washington, D.C., 1963).
- [159] Shinn, A. F., Food Crops and Postattack Recovery, Proc. Symp. on Post-attack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 23-42 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April 1968).
- [160] Osburn, W. S., Forecasting Long-Range Recovery from Nuclear Attack, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 107-136 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April 1968).
- [161] Brown, S., and Pilz, U., U.S. Agriculture: Potential Vulnerabilities, Report for DCPA, 83 pages (Stanford Research Institute, Menlo Park, Calif., Jan. 1969).

- [162] Comar, C. L., Wentworth, R. A., and Lengeman, F. W., Incorporation and Control of Fission Products Ingested by Livestock, Report for OCD, 36 pages (Dept. of Physical Biology, N.Y. State Veterinary College, Cornell U., Ithaca, N.Y., Oct. 1970).
- [163] Comar, C. L., Wentworth, R. A., and Lengemann, F. W., Incorporation and Control of Strontium, Cesium, and Iodine Secretion in Milk, Report for OCD, 64 pages (Dept. of Physical Biology, N.Y. State Veterinary College, Cornell U., Ithaca, N.Y., July 1969).
- [164] Protection of the Thyroid Gland in the Event of Releases of Radioiodine, NCRP Report No. 55, 60 pages (Natl. Council on Radiation Protection and Measurements, Washington, D.C. 20014, August 1, 1977).
- [165] Effects of Fallout on Agricultural Ecosystems in California, Report for OCD, 90 pages (U. of Calif., Riverside, Calif., May 1972).
- [166] Insect Pests of Major Food Crops, Their Reinvasion Potential and the Effects of Radiation on Arthropods, Report No. TRC-68-70, 199 pages (U. of Calif., Riverside, Calif., June 1969).
- [167] Auerback, S. I., Postattack Insect Problems, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 137-142 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April, 1968).
- [168] Reitemeier, R. F., Soil-Plant Relationships of Radioactive Nuclides in Fallout, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 143-148 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April, 1968).
- [169] Tompkins, P. C., Rise and Fall of Important Fission Products in the United States Food Supply, Proc. Symp. on Postattack Recovery from Nuclear War, Nov. 6-9, 1967, Ft. Monroe, Va., 169-170 (Office of Civil Defense or Natl. Acad. of Sciences, Washington, D.C., April 1968).
- [170] Blincoe, C., and Bohman, V. R., Decrease of Bovine Cesium-137 Concentrations Following the Cessations of Atmospheric Nuclear Weapons Testing, Advances in Chemistry Series 93, 427-435 (American Chemical Society, Washington, D.C. 1970).
- [171] Russell, R. S., Bartlett, B. O., and Bruce, R. S., The Significance of Long-Lived Nuclides after a Nuclear War, AEC Symposium Series 24, 548-565 (December 1971).



VIII.1 Comparison of Experimental and Calculated Exposures from Fallout Radiation as a Function of Depth in a Concrete Medium.
 O, X Experimental, Titus [1]; — Calculated $L(X)$ ($\times 10$) (Fig. V.12).

REFERENCES

- [1] F. Titus, "Penetration in Concrete of Gamma Radiation from Fallout," NBS Report 6143, (Sept. 1958).
- [2] C. McDonnell and J. Velletri, "An Experimental Evaluation of Roof Reduction Factors," CONESCO Report PSDC-TR-16, Flow Corporation, May 1966.
- [3] C. McDonnell, J. V. Velletri, A. W. Starbird, and J. F. Batter, "The Barrier Attenuation Introduced by a Vertical Wall," CONESCO Report PSDC-TR-15, Flow Corporation, September 1964.
- [4] R. M. Rubin, R. E. Faw, M. J. Coolbaugh, and J. M. Royer, "Energy and Angular Distributions of ⁶⁰Co Gamma Rays Penetrating a Concrete Shield," Nuc Eng and Des 16, 429, 1971. See Also "R. E. Faw, R. M. Rubin, J. M. Royer, and M. J. Coolbaugh, "Measurement of Angular and Energy Spectra of Wall Transmitted Radiation" Special Report Number 92 (NRDL TRC-69-10) Kansas State University, (July 1969).
- [5] C. Eisenhauer and A. B. Chilton, "Angular Distribution of Scattered Gamma Rays from a Fan Source", Trans. Am. Nuc Soc. 14, 1, June 1971.
- [6] L. V. Spencer, "Structure Shielding Against Fallout Radiation from Nuclear Weapons," NBS Monograph 42, National Bureau of Standards, June 1962.
- [7] Z. G. Burson and R. L. Summers, "Barrier Attenuation of Air-Scattered Gamma Radiation", CEX-63.3, Atomic Energy Commission, June 1965.
- [8] A. L. Kaplan, "Analysis of Data from Structure Shielding Experiments", NRDL-TRC-68-52, Technical Operations, Inc., June 1968.
- [9] M. A. Hughes and F. A. Bryan, Jr., "Effect of Source Field Elongation on Radiation Received at a Detector", Research Memorandum RM-333-4, Research Triangle Institute, Research Triangle Park, N.C., June 1968.
- [10] M. A. Schmoke and R. E. Rexroad, "Attenuation of Simulated Fallout Radiation by the Roof of a Concrete Blockhouse", Nuclear Defense Laboratory NDL-TR-6, August 1961.
- [11] M. A. Schmoke and R. E. Rexroad, "Attenuation of Fallout Radiation as a Function of Concrete Blockhouse Wall Thickness", NDL-TR-43, Nuclear Defense Laboratory, Oct. 1963.
- [12] R. Spring and C. H. McDonnell, "An Experimental Evaluation of Roof Reduction Factors within a Multistory Structure," CONESCO Report PSDC-TR-16, Supplement No. 1, April 1967.
- [13] C. McDonnell and J. Velletri, "Radiation Distribution within a Multi-story Structure", CONESCO Report TR-24, Flow Corporation, Feb. 1967.
- [14] A. B. Chilton, "Radiation Shielding, Analysis and Design Principles as Applied to Nuclear Defense Planning, Vol. IV Experimental Program," OCD-KSU Report Tr-40, W. R. Kimel, ed., Nov. 1966).